

Association Between Optimism and Serum Antioxidants in the Midlife in the United States Study

JULIA K. BOEHM, PhD, DAVID R. WILLIAMS, PhD, ERIC B. RIMM, ScD, CAROL RYFF, PhD, AND LAURA D. KUBZANSKY, PhD

Objective: Psychological and physical health are often conceptualized as the absence of disease; there is less research that addresses positive psychological and physical functioning. For example, optimism has been linked with reduced disease risk and biological dysfunction, but very little research has evaluated associations with markers of healthy biological functioning. Thus, we investigated the association between two indicators of positive health: optimism and serum antioxidants. **Methods:** The cross-sectional association between optimism and antioxidant concentrations was evaluated in 982 men and women from the Midlife in the United States study. Primary measures included self-reported optimism (assessed with the revised Life Orientation Test) and serum concentrations of nine different antioxidants (carotenoids and vitamin E). Regression analyses evaluated the relationship between optimism and antioxidant concentrations in models adjusted for demographics, health status, and health behaviors. **Results:** For every standard deviation increase in optimism, carotenoid concentrations increased by 3% to 13% in age-adjusted models. Controlling for demographic characteristics and health status attenuated this association. Fruit and vegetable consumption and smoking status were identified as potential pathways underlying the association between optimism and serum carotenoids. Optimism was not significantly associated with vitamin E. **Conclusions:** Optimism was associated with greater carotenoid concentrations, and this association was partially explained by diet and smoking status. The direction of effects cannot be conclusively determined. Effects may be bidirectional given that optimists are likely to engage in health behaviors associated with more serum antioxidants, and more serum antioxidants are likely associated with better physical health that enhances optimism. **Key words:** optimism, antioxidants, carotenoids, vitamin E.

GEE = generalized estimating equation; M = mean; MIDUS = Midlife in the United States; SD = standard deviation; SE = standard error.

INTRODUCTION

Although the World Health Organization defines health as “a state of... well-being and not merely the absence of disease or infirmity” (1), most empirical research has characterized both psychological and physical health by the presence or absence of disease. For example, psychological health has typically been assessed with indicators such as depression and anxiety, whereas physical health has been assessed with indicators such as hypertension and cardiovascular disease. However, health can only be understood comprehensively if the full spectrum of human functioning is considered. Therefore, characterizations of health need to include markers of enhanced psychological and

biological processes that indicate the presence of well-being, rather than only its absence. Indeed, researchers are increasingly recognizing that psychological well-being (including constructs such as optimism, happiness, and life purpose) is not the direct opposite of psychological ill-being, and that psychological well-being may have unique effects separate from those of psychological ill-being (2,3). Thus far, research that defines physical well-being in the context of positive biological functioning remains scarce.

Antioxidants, such as carotenoids (e.g., β-carotene) and vitamin E (e.g., α-tocopherol), constitute well-recognized markers of biological health. Antioxidants help to inhibit other molecules from oxidizing or producing free radicals that damage cells and contribute to disease pathophysiology (4). Dietary antioxidants are often assessed through self-reported inventories of food consumption and have been linked with reduced risk of cardiovascular disease (5,6). Objectively assessed serum antioxidants are also inversely associated with inflammation, atherosclerosis, diabetes, hypertension, coronary heart disease, and cancer (7–12). Although a substantial amount of work has evaluated how diet is associated with psychosocial factors (e.g., 13–15), less work has considered the association between serum antioxidants and psychosocial factors. To date, inverse associations have been reported for serum antioxidants and psychosocial factors such as socioeconomic status (16–18), poor cognitive function (19), depression (20,21), hostility (22), and job stress (23). However, no empirical research has investigated the association between psychological well-being and serum antioxidants.

Therefore, we investigated the cross-sectional association between optimism—an indicator of adaptive psychological functioning characterized by positive expectations for the future (24)—and serum carotenoids and vitamin E. We hypothesized that more optimistic individuals within a sample of aging adults would have greater concentrations of antioxidants than less optimistic individuals. We also investigated whether the association between optimism and antioxidant status was altered by potential confounders (i.e., demographic characteristics and health status).

From the Departments of Society, Human Development, and Health (J.K.B., D.R.W., L.D.K.), Nutrition (E.B.R.), and Epidemiology (E.B.R.), Harvard School of Public Health, Boston, Massachusetts; Department of African and African American Studies and Sociology (D.R.W.), Harvard University, Cambridge, Massachusetts; Channing Laboratory (E.B.R.), Department of Medicine, Brigham and Women’s Hospital and Harvard Medical School, Boston, Massachusetts; and Department of Psychology (C.R.) and Institute on Aging (C.R.), University of Wisconsin-Madison, Madison, Wisconsin.

Address correspondence and reprint requests to Julia K. Boehm, PhD, Department of Society, Human Development, and Health, Harvard School of Public Health, 677 Huntington Avenue, Boston, MA 02115. E-mail: jboehm@hsph.harvard.edu

Support for this research was provided by the Robert Wood Johnson Foundation through a grant, “Exploring the Concept of Positive Health,” to the Positive Psychology Center of the University of Pennsylvania, Martin Seligman, project director. The original Midlife in the United States study was supported by The John D. and Catherine T. MacArthur Foundation Research Network on Successful Midlife Development. Follow-up data collection was supported by the National Institute on Aging (P01-AG020166).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal’s Web site (www.psychosomaticmedicine.org).

Received for publication February 27, 2012; revision received August 22, 2012.
DOI: 10.1097/PSY.0b013e31827c08a9

OPTIMISM AND SERUM ANTIOXIDANTS

In addition, because of the strong link between antioxidant status and health behaviors such as fruit and vegetable consumption (19,25,26) and multivitamin supplementation (27), we separately evaluated whether health behaviors play a role in the pathway linking optimism and antioxidant concentrations.

METHODS

Participants

Participants were men and women drawn from the Midlife in the United States (MIDUS) study, which was initiated to understand a range of factors that influence the mental and physical health of Americans as they age. The first phase of the MIDUS study took place in 1994 to 1995 and comprised a national sample of 7108 individuals aged 25 to 74 years. Participants were recruited via random digit dialing and included siblings and twin pairs for some individuals (28).

The second phase of the MIDUS study began 9 to 10 years later (2004–2005) and involved five distinct projects (psychosocial assessment, daily diary study, cognitive function assessment, biomarker assessment, and neuroscience assessment). The current investigation included a subset of participants from Phase 2 who completed both the survey assessments of psychosocial function and the biomarker project. Individuals healthy enough to travel to the research clinic and who had completed the Phase 2 psychosocial assessment ($n = 5895$) were eligible to participate in the biomarker project (29). Of the 3191 individuals who were eligible to participate in the biomarker project, 39.3% ($n = 1255$) did so (29). After accounting for those individuals who could not be contacted, the response rate for the biomarker project was 43.1% (29). Notably, past work has demonstrated that participants in the biomarker project did not differ from participants in the psychosocial assessment by age, sex, race, marital status, income, chronic conditions, or body mass index; however, participants in the biomarker project were more highly educated than participants in the psychosocial assessment (29).

The analytic sample for the present study comprised 982 individuals with complete data on optimism, antioxidants, and covariates (of those, 365 were members of a sibling or twin set). Participants provided informed consent, and the research was approved by institutional review boards at participating institutions.

Measures

Because the current investigation combined data from two separate projects that took place during Phase 2 of the MIDUS study, not all variables were assessed at the same point in time. Optimism and demographic characteristics were assessed before the measurement of antioxidants, health status, and health behaviors. The latter set of variables was measured, on average, 830.62 days (a little more than 2 years) after optimism and the demographic characteristics had been measured (standard deviation [SD] = 451.71; range, 115–1893 days).

Optimism

Optimism (i.e., the general expectation that the future will be favorable) was assessed with the well-validated six-item revised Life Orientation Test (30). Sample items include the positively worded "I expect more good things to happen to me than bad" and the negatively worded "I hardly ever expect things to go my way." Participants responded to each item with the following scale: 1 = agree a lot, 2 = agree a little, 3 = neither agree nor disagree, 4 = disagree a little, and 5 = disagree a lot. The total optimism score was computed by first reverse-scoring the three positively worded items, summing the responses of all six items, and finally standardizing the score (mean [M] = 0, SD = 1). Larger scores indicated higher levels of optimism, and internal consistency reliability was good in this sample ($\alpha = 0.81$). Prior work with other participants has reported comparable internal consistency reliabilities (e.g., $\alpha = 0.79$) and test-retest reliabilities ranging from 0.56 to 0.79 across 4 to 28 months (30). The revised Life Orientation Test has also previously demonstrated acceptable convergent and discriminant validity (30).

According to MIDUS study data conventions, missing values were imputed from the mean of completed items if at least three of the six items were available. In addition to a continuous measure of optimism, we assessed potential

discontinuities in effects by creating tertiles based on the distribution of scores. Standardized scores of -0.40 or lower were categorized as low (33.95%), scores higher than -0.40 and lower than 0.50 were categorized as moderate (30.27%), and scores of 0.50 or higher were categorized as high (35.79%).

Although researchers have sometimes split the revised Life Orientation Test into subscales based on the valence of items (positively worded versus negatively worded), we opted not to do this for several reasons. It is increasingly apparent that this practice may be at odds with the goal of controlling for acquiescence response bias in the measurement of psychological constructs. Moreover, a comprehensive assessment of optimism requires that respondents not only endorse positively worded items but also disagree with negatively worded items (31). Finally, valence-based subscales compromised the internal consistency of the optimism scale ($\alpha_{\text{optimism subscale}} = 0.66$; $\alpha_{\text{pessimism subscale}} = 0.82$).

Antioxidants

After traveling to one of three clinical research centers (University of Wisconsin, Georgetown University, and University of California-Los Angeles) for a 2-day visit, participants provided a fasting blood sample on the second morning of their visit. Blood samples were processed with standardized procedures at each research center and stored in a freezer between -60°C and -80°C . Assays of the frozen samples were conducted at the Antioxidants Research Laboratory at Tufts University. In accordance with previous methodology (31,32), a reverse-phase gradient high-pressure liquid chromatography system was used with ultraviolet-visible and fluorescence detection.

Measures of the following nine antioxidants (μM) were obtained: carotenoids included *trans*- β -carotene, 13-*cis*- β -carotene, α -carotene, β -cryptoxanthin, lutein, zeaxanthin, and lycopene; vitamin E included α -tocopherol and γ -tocopherol. We created a composite of total carotenoids that was composed of *trans*- β -carotene, α -carotene, β -cryptoxanthin, lutein, and zeaxanthin, but excluded lycopene because it has demonstrated inconsistent associations with health-related variables (12,16,22). Serum concentrations of each carotenoid were first standardized ($M = 0$, $SD = 1$) and then summed to represent total carotenoid levels. One outlier was removed to maintain a normal distribution for the carotenoid composite, yielding a sample size of 981 in relevant statistical analyses.

For carotenoids, the minimal assay range was 0.01 to 0.03 μM , the interassay coefficient of variation ranged from 1% to 16%, and the intra-assay coefficient of variation ranged from 6% to 13%. The minimum assay range was 1.18 μM for α -tocopherol and 0.11 μM for γ -tocopherol. The interassay coefficient of variation for the tocopherols ranged from 2% to 6%, and the intra-assay coefficient of variation ranged from 6% to 10%.

Potential Confounders

Self-reported confounders included demographic characteristics and health status. Demographic factors were age (standardized years), sex (male, female), race/ethnicity (white, nonwhite), education (less than high school, high school degree, some college, 4-year college degree or higher), and household income (standardized US\$). Health status factors included chronic conditions (heart disease, hypertension, stroke, high cholesterol, diabetes, or cancer [none, presence of one or more]) and body mass index (standardized kg/m^2). Categorical variables were dummy-coded and continuous variables were standardized ($M = 0$, $SD = 1$) before inclusion in the models.

Potential Pathways

Health behaviors (fruit and vegetable consumption, multivitamin supplementation, exercise, and smoking status) were investigated as possible pathways by which optimism and antioxidant status may be associated.¹ Participants reported the mean number of fruit and vegetable servings they consumed in a typical day (two or fewer, three to four, five or more) and their consumption of multivitamin supplements (no, yes). Participants also indicated whether they

¹Alcohol consumption was also considered as a potential pathway variable but made no difference in regression models and will not be discussed further. In addition, findings that controlled for depressed affect are discussed in the supplementary text (Supplemental Digital Content 1, <http://links.lww.com/PSYMED/A59>).

TABLE 1. Distribution of Participant Characteristics by Level of Optimism

Characteristics	Optimism			<i>p</i>
	Low (<i>n</i> = 332)	Moderate (<i>n</i> = 298)	High (<i>n</i> = 352)	
Age, M (SD), y	53.43 (11.71)	56.55 (12.32)	56.01 (11.28)	.002
Sex, %				
Men	48.49	44.97	42.33	.27
Women	51.51	55.03	57.67	
Race, %				
White	91.57	94.97	93.47	.23
Nonwhite	8.43	5.03	6.53	
Education, %				
Less than high school	5.42	2.35	2.56	<.001
High school degree	26.51	20.81	15.06	
Some college	32.23	29.87	25.85	
4-y college degree or higher	35.84	46.98	56.53	
Household income, M (SD), US\$	68,006.96 (54,411.13)	75,354.99 (58,081.43)	84,756.40 (64,189.53)	.001
Time between assessments, M (SD), d	828.87 (452.58)	831.31 (452.71)	831.69 (451.33)	.996
Chronic conditions, % ^a				
Yes	64.16	64.77	63.07	.90
No	35.84	35.23	36.93	
Body mass index, M (SD), kg/m ²	29.74 (6.48)	28.73 (5.51)	28.78 (5.67)	.049
Fruit and vegetable consumption, %				
Two or fewer servings per day	49.70	39.93	32.95	<.001
Three to four servings per day	34.04	37.58	42.33	
Five or more servings per day	16.27	22.48	24.72	
Multivitamin supplementation, %				
Yes	51.20	55.70	52.56	.51
No	48.80	44.30	47.44	
Regular exercise, %				
Yes	74.10	83.89	80.11	.009
No	25.90	16.11	19.89	
Smoking status, %				
Current smoker	17.77	8.72	6.82	<.001
Past smoker	30.72	36.58	32.67	
Never smoker	51.51	54.70	60.51	

M = mean; SD = standard deviation.

Percentages refer to the column percentage of individuals within each optimism category who have that characteristic. *P* value comes from χ^2 test or analysis of variance.

^a Because our measure of chronic conditions included hypertension and high cholesterol, the proportion of participants with chronic conditions may seem relatively large; however, the prevalence of such conditions in our sample is consistent with the prevalence of such conditions among middle-aged to older adults living in the United States (34).

had engaged in regular exercise at least three times each week for 20 minutes (no, yes) and their smoking status (current smoker, past smoker, never smoker). All behavior-related variables were dummy coded.

Statistical Analyses

Statistical analyses were conducted in SAS version 9.2. Individual antioxidant concentrations were log-transformed (base e) before inclusion in the analyses to improve distributional properties.² The log-transformed variables had approximately normal distributions. For each antioxidant and the total

carotenoid score, the primary analyses included a series of hierarchical linear regressions with the continuous optimism score. We first fitted age-adjusted models with optimism as a predictor of each antioxidant concentration. A second minimally adjusted model contained all demographic characteristics (age, sex, race/ethnicity, education, and income). Because the amount of time between the assessment of optimism and the assessment of serum antioxidants differed for each participant, we also included a covariate to account for this in the minimally adjusted model. Although optimism and antioxidant levels are considered to be largely stable over time (30,33), instability in either measure would likely bias any observed associations toward the null; thus, these analyses may provide a conservative test of our hypothesis. The third model added health status (chronic conditions and body mass index) to the second model.

Additional models tested whether health behaviors independently accounted for any part of the association between optimism and antioxidant concentrations in models minimally adjusted for demographics (age, sex, race, education, income,

²One participant with a 13-cis-β-carotene concentration of 0 and one participant with a zeaxanthin concentration of 0 were set as missing because the values were either an aberrant finding or below the detection level of the methodology.

OPTIMISM AND SERUM ANTIOXIDANTS

and time between assessments). To this end, we compared the regression coefficients for optimism when a potential pathway variable was either excluded from the model or included in the model. When the regression coefficient for optimism was reduced with the inclusion of a potential pathway variable (as indicated by the percentage change), this was taken to suggest that the potential pathway variable accounted for part of the association between optimism and antioxidant concentrations.

Secondary analyses tested for potential discontinuities in effects by including tertiles of optimism in models that were age-adjusted and minimally adjusted. We then repeated the primary statistical analyses with generalized estimating equations (GEEs) to account for the data clustering of the sibling and twin sets. Results were virtually unchanged in GEE analyses, suggesting that clustering of families did not bias parameter estimates or standard errors. Findings from non-GEE analyses are presented in the text for ease of interpretation, but see Supplementary Tables 1 and 2 (<http://links.lww.com/PSYMED/A60>, <http://links.lww.com/PSYMED/A61>) for the primary analyses with GEE.

RESULTS

Sample Characteristics

The analytic sample was composed of men (45.21%) and women (54.79%) with a M (SD) age of 55.30 (11.81) years (range, 34–84 years). Most participants were white (93.28%), and 46.64% had a 4-year college degree or higher (29.23% had some college, 20.67% had a high school degree, and 3.46% had less than a high school degree). Table 1 shows sample characteristics according to the level of optimism. Individuals with low optimism levels were significantly younger and reported less income and education than those with higher optimism levels. In addition, less optimistic individuals tended to be current smokers,

to engage in less regular exercise, and to consume fewer fruits and vegetables.

The unadjusted M (SD) for each antioxidant was as follows (in μM): *trans*- β -carotene, 0.63 (0.75); 13-*cis*- β -carotene, 0.08 (0.08); α -carotene, 0.10 (0.11); β -cryptoxanthin, 0.21 (0.18); lutein, 0.30 (0.21); zeaxanthin, 0.07 (0.05); lycopene, 0.46 (0.23); α -tocopherol, 29.36 (13.67); γ -tocopherol, 3.48 (2.73). The M (SD) total carotenoid composite of the individual standardized carotenoids was -0.03 (3.12).

Association Between Optimism and Antioxidants

Individuals with greater optimism were generally more likely to have higher concentrations of carotenoids (Table 2); that is, for *trans*- β -carotene, α -carotene, β -cryptoxanthin, lutein, zeaxanthin, and lycopene, age-adjusted and minimally adjusted regression models indicated that every standard deviation increase in optimism was associated with 3% to 13% higher carotenoid concentrations. Although these findings were sometimes attenuated to marginal statistical significance or nonsignificance with the addition of health status, the magnitude of the regression coefficients remained similar, and 3% to 7% higher carotenoid concentrations were still evident. Findings for the total carotenoid composite had a similar pattern in age-adjusted models, minimally adjusted models, and models adjusted for health status (Table 2). Namely, optimism was positively and significantly associated with total carotenoid concentrations. Optimism was not significantly associated with α -tocopherol and γ -tocopherol. Given

TABLE 2. Nonstandardized Parameter Estimates (Standard Error) for the Association Between Every Standard Deviation Increase in Optimism and Serum Antioxidants ($N = 982$)

Log Antioxidants, μM^a	Optimism		
	Model 1 ^b	Model 2 ^c	Model 3 ^d
Carotenoids			
<i>trans</i> - β -Carotene	0.10*** (0.03)	0.06* (0.03)	0.04 (0.03)
13- <i>cis</i> - β -Carotene	0.02 (0.02)	-0.004 (0.02)	-0.01 (0.02)
α -Carotene	0.13*** (0.03)	0.08** (0.03)	0.07* (0.03)
β -Cryptoxanthin	0.07*** (0.02)	0.05* (0.02)	0.04* (0.02)
Lutein	0.05** (0.02)	0.04* (0.02)	0.03 [†] (0.02)
Zeaxanthin	0.04* (0.02)	0.03 [†] (0.02)	0.03 (0.02)
Lycopene	0.03 [†] (0.02)	0.04* (0.02)	0.04* (0.02)
Total carotenoids	0.39*** (0.10)	0.24* (0.10)	0.19* (0.10)
Vitamin E			
α -Tocopherol	0.001 (0.02)	-0.002 (0.02)	0.0003 (0.02)
γ -Tocopherol	-0.02 (0.02)	-0.007 (0.02)	-0.003 (0.02)

[†] $p \leq .10$.

^{*} $p \leq .05$.

^{**} $p \leq .01$.

^{***} $p \leq .001$.

^a Each individual antioxidant was log-transformed; the composite of the total carotenoids was not.

^b Adjusted for age.

^c Adjusted for demographics (age, sex, race, education, income, and time between assessments).

^d Adjusted for demographics and health status (chronic conditions and body mass index).

Bold values signify statistical significance.

optimism's null association with tocopherols in these and subsequent analyses, we will not discuss results for vitamin E further.

Health Behaviors as Potential Pathways Linking Optimism and Total Carotenoids

Given that carotenoid concentrations demonstrated the most robust association with optimism, we investigated whether the association was potentially accounted for by health behaviors. To start, we evaluated the association between the total carotenoid composite and each health behavior. Individuals who consumed five or more servings of fruits and vegetables each day had higher carotenoid concentrations ($M = 1.25$, $SD = 3.40$) relative to those who consumed three or four servings ($M = 0.17$, $SD = 3.02$) or two or fewer servings ($M = -0.89$, $SD = 2.79$), $F(2,978) = 34.77$, $p < .001$. Similarly, individuals who consumed multivitamins had higher carotenoid concentrations ($M = 0.30$, $SD = 3.06$) than those who did not ($M = -0.40$, $SD = 3.15$), $t(979) = 3.51$, $p < .001$. Individuals who exercised had higher carotenoid concentrations ($M = 0.14$, $SD = 3.17$) than those who did not ($M = -0.70$, $SD = 2.85$), $t(979) = 3.44$, $p < .001$. Current smokers had the lowest carotenoid concentrations ($M = -1.60$, $SD = 2.30$) relative to past smokers ($M = -0.006$, $SD = 3.10$) and never smokers ($M = 0.26$, $SD = 3.18$), $F(2,978) = 16.77$, $p < .001$.

We next determined how the association between optimism and total carotenoids changed when controlling for each poten-

tial pathway variable and demographic characteristics. As shown in Table 3, fruit and vegetable consumption substantially reduced the association between optimism and total carotenoids, as did smoking status. In both cases, the initially statistically significant regression coefficient for optimism was reduced by 25% or more and became marginally significant when the potential pathway variable was included. Exercise also explained part of the association, but multivitamin supplementation did not. All of the health behaviors combined accounted for nearly half of the association between optimism and total carotenoid concentrations and reduced the association to nonsignificance.

Secondary Analyses

Findings for tertiles of optimism were consistent with those described previously (Table 4). Many of the same carotenoids that were associated with optimism continuously were associated with high levels of optimism in age-adjusted models and, to a lesser extent, in minimally adjusted models. That is, relative to the lowest levels of optimism, the highest levels of optimism were associated with 9% to 20% higher concentrations of *trans*- β -carotene, α -carotene, β -cryptoxanthin, lutein, zeaxanthin, and lycopene in age-adjusted models. In addition, for the total carotenoid composite, the highest level of optimism was significantly associated with greater total carotenoids in age-adjusted and minimally adjusted models. Thus, compared with the lowest optimism levels, the highest levels were positively associated with

TABLE 3. Change in the Association Between Optimism and Total Carotenoid Concentrations When Potential Pathway Variables Are Included ($N = 981$)

Predictors	Association With Total Carotenoid Concentrations ^a		
	<i>b</i>	SE	% Δ
Optimism			
Demographic covariates ^b	0.24*	0.10	—
Optimism			
Demographic covariates ^b	0.17 [†]	0.10	-29
Fruit and vegetable consumption			
Optimism			
Demographic covariates ^b	0.24*	0.10	0
Multivitamin supplementation			
Optimism			
Demographic covariates ^b	0.22*	0.10	-8
Exercise			
Optimism			
Demographic covariates ^b	0.18 [†]	0.10	-25
Smoking status			
Optimism			
Demographic covariates ^b	0.13	0.10	-46
All behaviors ^c			

SE = standard error.

p values symbolized in the second column from the left indicate the association between optimism and total carotenoid concentrations in the presence of demographic covariates or demographic covariates plus specified pathway variables.

[†] $p \leq .10$.

* $p \leq .05$.

^a Total carotenoid concentrations were composed of *trans*- β -carotene, α -carotene, β -cryptoxanthin, lutein, and zeaxanthin.

^b Adjusted for age, sex, race, education, income, and time between assessments.

^c Adjusted for fruit and vegetable consumption, multivitamin supplementation, exercise, and smoking status.

OPTIMISM AND SERUM ANTIOXIDANTS

TABLE 4. Nonstandardized Parameter Estimates (Standard Error) for the Association Between Tertiled Optimism and Serum Carotenoids in Age-Adjusted and Minimally Adjusted Models^a (*N* = 982)^b

Log Carotenoids, μM^c	Optimism		
	Low (<i>n</i> = 332)	Moderate (<i>n</i> = 298)	High (<i>n</i> = 352)
<i>trans</i> - β -Carotene			
Age-adjusted	Reference	0.08 (0.07)	0.20** (0.07)
Minimally adjusted	Reference	0.02 (0.07)	0.11 (0.07)
13- <i>cis</i> - β -Carotene			
Age-adjusted	Reference	-0.008 (0.06)	0.07 (0.06)
Minimally adjusted	Reference	-0.05 (0.06)	-0.003 (0.06)
α -Carotene			
Age-adjusted	Reference	0.14 [†] (0.08)	0.20** (0.07)
Minimally adjusted	Reference	0.07 (0.08)	0.10 (0.07)
β -Cryptoxanthin			
Age-adjusted	Reference	0.10* (0.05)	0.17*** (0.05)
Minimally adjusted	Reference	0.09 [†] (0.05)	0.14** (0.05)
Lutein			
Age-adjusted	Reference	0.04 (0.05)	0.12** (0.05)
Minimally adjusted	Reference	0.03 (0.04)	0.09* (0.04)
Zeaxanthin			
Age-adjusted	Reference	0.04 (0.05)	0.10* (0.04)
Minimally adjusted	Reference	0.03 (0.05)	0.08 [†] (0.04)
Lycopene			
Age-adjusted	Reference	0.03 (0.04)	0.09* (0.04)
Minimally adjusted	Reference	0.04 (0.04)	0.10** (0.04)
Total carotenoids			
Age-adjusted	Reference	0.37 (0.25)	0.85*** (0.24)
Minimally adjusted	Reference	0.19 (0.24)	0.51* (0.24)

[†] $p \leq .10$.

* $p \leq .05$.

** $p \leq .01$.

*** $p \leq .001$.

^a Adjusted for demographics (age, sex, race, education, income, and time between assessments).

^b Analyses for total carotenoids were based on *N* = 981 owing to one excluded outlier.

^c Each individual carotenoid was log-transformed; the composite of the total carotenoids was not.

carotenoid concentrations. Evidence for an association between moderate and low levels was less strong. However, because statistical power is reduced in tertiled analyses, these findings warrant cautious interpretation.

DISCUSSION

This is the first study to investigate the association between self-reported optimism and serum antioxidants. Whereas previous health research has mostly emphasized problematic psychological and physical functioning (i.e., the presence of ill-being or disease), we used a positive definition of health by focusing on psychological and biological processes that indicate the presence of restorative processes rather than the absence of deteriorative processes. Findings suggest that optimism is positively associated with greater concentrations of one group of antioxidants, namely, carotenoids. This relationship held for both continuous and categorical measures of optimism. The relationship also remained mostly statistically significant after controlling

for demographic characteristics (e.g., socioeconomic status) and health status (e.g., body mass index), although effect sizes were somewhat attenuated with the addition of these factors. Although optimism was not associated with vitamin E, previous research has also reported null associations between psychosocial factors and tocopherols (22,35). Tocopherols may be more strongly dependent on other factors (e.g., lipids), which may make it difficult to detect the direct effects of psychosocial variables.

One of the major strengths of the current study is its objective assessment of serum antioxidants, which alleviates concerns that optimism's observed associations are caused by reporting bias. Moreover, including health behaviors individually in regression models allowed the investigation of the potential pathways linking optimism and carotenoids. The role of health behaviors (in particular, diet and vitamin consumption) has rarely been explored in previous research. Findings indicated that the consumption of fruits and vegetables, along with smoking status,

explained the greatest variation in the association between optimism and total carotenoid concentrations. Indeed, individuals who ate two or fewer servings of fruits and vegetables a day were significantly less optimistic than individuals who ate three or more servings a day. These findings are consistent with research suggesting that optimism is linked with consuming a healthy diet, exercising regularly, and not smoking cigarettes (36–39). The favorable expectations that optimistic individuals hold for the future may provide them with the resources to better manage challenge and stress (40), particularly in the context of health behaviors. Although most people want to lead a healthy life-style, daily constraints often thwart such intentions. Because optimists tend to persist at their goals, use successful coping strategies, and know when to pursue achievable versus unachievable goals (40), they may be better able to maintain the healthy behaviors that are associated with improved antioxidant status. Of course, the reverse may also be true; that is, individuals who engage in healthier behaviors and enjoy better health may also be more optimistic about life.

Although this study confirms that the association between optimism and serum antioxidants can be partially explained by fruit and vegetable consumption and cigarette smoking, it is also interesting to note that including health behaviors did not completely explain all of the variance in the relationship. Although this may suggest that measurement error was present in the assessment of self-reported health behaviors, we note that these are commonly used measures and that including them is a unique strength of the study. Thus, assuming there is limited residual confounding, such findings could also indicate that other pathways may be operating. For example, one possibility is that optimism and antioxidant concentrations are associated because positive psychological factors enhance the bodily processes that encourage antioxidant absorption. Although the present research cannot address this issue, early studies on positive psychological states, such as love and comfort, suggest that digestion may vary depending on the psychological context (41). Moreover, recent evidence indicates that numerous factors influence the bioavailability of antioxidants and that patterns of bioavailability can vary by type of antioxidant (42,43). Another possibility comes from work suggesting that levels of serum antioxidants are lower in the presence of greater inflammation (12,44). In fact, this explanation has been proposed for why smoking is associated with reduced antioxidant levels (45,46). Given prior work suggesting that higher levels of optimism are associated with reduced inflammation (47,48), mitigating inflammatory processes may be another pathway by which optimism is associated with higher antioxidant concentrations. Thus, it is feasible that optimism and serum antioxidants may be linked by as yet unidentified physiological processes.

Evidence that antioxidant supplementation alone does not improve disease or mortality outcomes is also worth noting (e.g., 49). Although a conservative interpretation of this evidence might suggest that serum antioxidants serve primarily as a marker of health habits, prior research has not ruled out the possibility that antioxidant supplementation may have limited effects on health outcomes because effects are influenced by interactions with

other biological substrates. Thus, antioxidants may yet emerge as a marker of biological health rather than merely a marker of healthy behaviors such as eating a balanced diet.

One of the major limitations of the current study is its cross-sectional design. This prevents conclusions about the causal direction of the optimism-antioxidant association. We suspect that it is likely bidirectional (i.e., optimistic individuals are likely to engage in health behaviors associated with greater concentrations of antioxidants; greater concentrations of antioxidants are associated with better physical health that may enhance optimism), but the time between assessments reduces the likelihood that serum antioxidants were driving optimism levels. Furthermore, health behaviors may confound the association between optimism and carotenoids rather than be on the pathway; however, the current data cannot resolve such issues. It is also possible that other unevaluated third variables could explain the association between optimism and antioxidant concentrations. For example, some evidence suggests that different concentrations of antioxidants may be associated with genetic variation (50), and levels of optimism could similarly have a genetic basis.

In addition, there was an approximately 2-year lag between the assessment of optimism and the assessment of serum antioxidants. Although concurrent assessment would have been preferable, both optimism and serum antioxidant measurements are relatively stable within periods of several years (30,33). Antioxidant concentrations were also measured on a single occasion, which may be susceptible to within-person variability. Future research with repeated measures of antioxidant concentrations may provide a better understanding of typical concentrations, although research also suggests that single assessments of antioxidant status reliably characterize longer-term concentrations (33). Another caveat concerns the use of a predominantly white, middle-aged sample, which limits the generalizability of the findings. We cannot be certain that the association between optimism and carotenoids is present in other ethnic groups or even at other stages of the life span.

Despite these limitations, this study is one of the first of its kind to investigate how positive indicators of psychological health (i.e., optimism) and physical health (i.e., serum antioxidants) might be associated. This study also adds to the relatively limited literature on psychosocial factors and serum antioxidant status. Additional strengths include a relatively large sample size, a comprehensive set of covariates, and objectively assessed antioxidant concentrations. Taken together, results suggest that optimism was associated with higher carotenoid concentrations. Moreover, health behaviors, such as fruit and vegetable consumption, seemed to explain a large proportion of the optimism-antioxidant relationship. Future research would benefit from the use of prospective study designs, diverse samples, and other measures that capture adaptive psychological functioning. Moreover, researchers could evaluate whether adaptive psychological functioning protects against challenges that are known to be associated with diminished antioxidant status such as racial inequality and low socioeconomic status. Additional research might evaluate whether manipulating or enhancing optimism could increase serum antioxidants or the likelihood of engaging in health behaviors. Replication of these

OPTIMISM AND SERUM ANTIOXIDANTS

findings and a more nuanced understanding of relevant pathways may indicate that enhancing psychosocial assets such as optimism could provide effective strategies for promoting health.

We thank the staff of the clinical research centers at Georgetown University, University of Wisconsin-Madison, and University of California-Los Angeles for their support in conducting this study.

REFERENCES

1. World Health Organization. WHO definition of health. In: Preamble to the Constitution of the World Health Organization, as adopted by the International Health Conference, New York, June 19 to July 22, 1946. Signed on July 22, 1946 by the representatives of 61 states (Official Records of the World Health Organization, no. 2, p. 100) and entered into force on April 7, 1948.
2. Ryff CD, Love GD, Urry HL, Muller D, Rosenkranz MA, Friedman EM, Davidson RJ, Singer B. Psychological well-being and ill-being: do they have distinct or mirrored biological correlates? *Psychother Psychosom* 2006;75:85–95.
3. Ryff CD, Singer B. The contours of positive human health. *Psycholog Inq* 1998;9:1–28.
4. Young IS, Woodside JV. Antioxidants in health and disease. *J Clin Pathol* 2001;54:176–86.
5. Knekut P, Ritz J, Pereira MA, O'Reilly EJ, Augustsson K, Fraser GE, Goldbourt U, Heitmann BL, Hallmans G, Liu S, Pietinen P, Spiegelman D, Stevens J, Virtamo J, Willett WC, Rimm EB, Ascherio A. Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. *Am J Clin Nutr* 2004;80:1508–20.
6. Rautiainen S, Larsson S, Virtamo J, Wolk A. Total antioxidant capacity of diet and risk of stroke: a population-based prospective cohort of women. *Stroke* 2012;43:335–40.
7. Giovannucci E. Tomatoes, tomato-based products, lycopene, and cancer: review of the epidemiologic literature. *J Natl Cancer Inst* 1999;91:317–31.
8. Coyne T, Ibiebele TI, Baade PD, Dobson A, McClinton C, Dunn S, Leonard D, Shaw J. Diabetes mellitus and serum carotenoids: findings of a population-based study in Queensland, Australia. *Am J Clin Nutr* 2005;82:685–93.
9. Hozawa A, Jacobs DR Jr, Steffes MW, Gross MD, Steffen LM, Lee DH. Circulating carotenoid concentrations and incident hypertension: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *J Hypertens* 2009;27:237–42.
10. Ribaya-Mercado JD, Blumberg JB. Lutein and zeaxanthin and their potential roles in disease prevention. *J Am Coll Nutr* 2004;23:567S–87S.
11. Gey KF, Moser UK, Jordan P, Stahelin HB, Eichholzer M, Ludin E. Increased risk of cardiovascular disease at suboptimal plasma concentrations of essential antioxidants: an epidemiological update with special attention to carotene and vitamin C. *Am J Clin Nutr* 1993;57:787S–97S.
12. Hozawa A, Jacobs DR Jr, Steffes MW, Gross MD, Steffen LM, Lee DH. Relationships of circulating carotenoid concentrations with several markers of inflammation, oxidative stress, and endothelial dysfunction: the Coronary Artery Risk Development in Young Adults (CARDIA)/Young Adult Longitudinal Trends in Antioxidants (YALTA) study. *Clin Chem* 2007;53:447–55.
13. Darmon N, Drewnowski A. Does social class predict diet quality? *Am J Clin Nutr* 2008;87:1107–17.
14. Sanchez-Villegas A, Delgado-Rodriguez M, Alonso A, Schlatter J, Lahortiga F, Majem LS, Martinez-Gonzalez MA. Association of the Mediterranean dietary pattern with the incidence of depression: the Seguimiento Universidad de Navarra/University of Navarra Follow-up (SUN) Cohort. *Arch Gen Psychiatry* 2009;66:1090–8.
15. Shaikh AR, Yaroch AL, Nebeling L, Yeh MC, Resnicow K. Psychosocial predictors of fruit and vegetable consumption in adults: a review of the literature. *Am J Prev Med* 2008;34:535–43.
16. Janicki-Deverts D, Cohen S, Matthews KA, Gross MD, Jacobs DR Jr. Socioeconomic status, antioxidant micronutrients, and correlates of oxidative damage: the Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Psychosom Med* 2009;71:541–8.
17. Nicklett EJ, Szanton S, Sun K, Ferrucci L, Fried LP, Guralnik JM, Semba RD. Neighborhood socioeconomic status is associated with serum carotenoid concentrations in older, community-dwelling women. *J Nutr* 2011;141:284–9.
18. Stimpson JP, Nash AC, Ju H, Eschbach K. Neighborhood deprivation is associated with lower levels of serum carotenoids among adults participating in the Third National Health and Nutrition Examination Survey. *J Am Diet Assoc* 2007;107:1895–902.
19. Polidori MC, Pratico D, Mangialasche F, Mariani E, Aust O, Anlasik T, Mang N, Pientka L, Stahl W, Sies H, Mecocci P, Nelles G. High fruit and vegetable intake is positively correlated with antioxidant status and cognitive performance in healthy subjects. *J Alzheimers Dis* 2009;17:921–7.
20. Maes M, De Vos N, Pioli R, Demedts P, Wauters A, Neels H, Christophe A. Lower serum vitamin E concentrations in major depression: another marker of lowered antioxidant defenses in that illness. *J Affect Disord* 2000;58:241–6.
21. Shibata H, Kumagai S, Watanabe S, Suzuki T. Relationship of serum cholesterol and vitamin E to depressive status in the elderly. *J Epidemiol* 1999;9:261–7.
22. Ohira T, Hozawa A, Iribarren C, Daviglus ML, Matthews KA, Gross MD, Jacobs DR Jr. Longitudinal association of serum carotenoids and tocopherols with hostility: the CARDIA Study. *Am J Epidemiol* 2008;167:42–50.
23. Tsuboi H, Tatsumi A, Yamamoto K, Kobayashi F, Shimoji K, Kinoshita N. Possible connections among job stress, depressive symptoms, lipid modulation and antioxidants. *J Affect Disord* 2006;91:63–70.
24. Carver CS, Scheier MF, Segerstrom SC. Optimism. *Clin Psychol Rev* 2010;30:879–89.
25. Block G, Norkus E, Hudes M, Mandel S, Helzlsouer K. Which plasma antioxidants are most related to fruit and vegetable consumption? *Am J Epidemiol* 2001;154:1113–8.
26. Campbell DR, Gross MD, Martini MC, Grandits GA, Slavin JL, Potter JD. Plasma carotenoids as biomarkers of vegetable and fruit intake. *Cancer Epidemiol Biomarkers Prev* 1994;3:493–500.
27. Wolters M, Hermann S, Hahn A. Effects of 6-month multivitamin supplementation on serum concentrations of alpha-tocopherol, beta-carotene, and vitamin C in healthy elderly women. *Int J Vitam Nutr Res* 2004;74:161–8.
28. Radler BT, Ryff CD. Who participates? Accounting for longitudinal retention in the MIDUS National Study of Health and Well-Being. *J Aging Health* 2010;22:307–31.
29. Dienberg Love G, Seeman TE, Weinstein M, Ryff CD. Bioindicators in the MIDUS National Study: protocol, measures, sample, and comparative context. *J Aging Health* 2010;22:1059–80.
30. Scheier MF, Carver CS, Bridges MW. Distinguishing optimism from neuroticism (and trait anxiety, self-mastery, and self-esteem): a reevaluation of the Life Orientation Test. *J Pers Soc Psychol* 1994;67:1063–78.
31. Ribaya-Mercado JD, Fox JG, Rosenblad WD, Blanco MC, Russell RM. Beta-carotene, retinol and retinyl ester concentrations in serum and selected tissues of ferrets fed beta-carotene. *J Nutr* 1992;122:1898–903.
32. Yeum KJ, Ahn SH, Rupp de Paiva SA, Lee-Kim YC, Krinsky NI, Russell RM. Correlation between carotenoid concentrations in serum and normal breast adipose tissue of women with benign breast tumor or breast cancer. *J Nutr* 1998;128:1920–6.
33. Kotopoulous J, Tworoger SS, Campos H, Chung FL, Clevenger CV, Franke AA, Mantzoros CS, Ricchiuti V, Willett WC, Hankinson SE, Eliassen AH. Reproducibility of plasma and urine biomarkers among premenopausal and postmenopausal women from the Nurses' Health Studies. *Cancer Epidemiol Biomarkers Prev* 2010;19:938–46.
34. Roger VL, Go AS, Lloyd-Jones DM, Benjamin EJ, Berry JD, Borden WB, Bravata DM, Dai S, Ford ES, Fox CS, Fullerton HJ, Gillespie C, Hailpern SM, Heit JA, Howard VJ, Kissela BM, Kittner SJ, Lackland DT, Lichtman JH, Lisabeth LD, Makuc DM, Marcus GM, Marelli A, Matchar DB, Moy CS, Mozaffarian D, Mussolini ME, Nichol G, Paynter NP, Soliman EZ, Sorlie PD, Sotoodehnia N, Turan TN, Virani SS, Wong ND, Woo D, Turner MB. American Heart Association Statistics Committee, Stroke Statistics Subcommittee. Heart disease and stroke statistics—2012 update: a report from the American Heart Association. *Circulation* 2012;125:e2–220.
35. Tiemeier H, Hofman A, Kiliaan AJ, Meijer J, Breteler MMB. Vitamin E and depressive symptoms are not related. The Rotterdam Study. *J Affect Disord* 2002;72:79–83.
36. Kavussanu M, McAuley E. Exercise and optimism: are highly active individuals more optimistic? *J Sport Exerc Psychol* 1995;17:246–58.
37. Giltay EJ, Geleijnse JM, Zitman FG, Buijsse B, Kromhout D. Lifestyle and dietary correlates of dispositional optimism in men: the Zutphen Elderly Study. *J Psychosom Res* 2007;63:483–90.
38. Kelloniemi H, Ek E, Laitinen J. Optimism, dietary habits, body mass index and smoking among young Finnish adults. *Appetite* 2005;45:169–76.
39. Steptoe A, Wright C, Kunz-Ebrecht SR, Iliffe S. Dispositional optimism and health behaviour in community-dwelling older people: associations with healthy ageing. *Br J Health Psychol* 2006;11:71–84.

40. Rasmussen HN, Wrosch C, Scheier MF, Carver CS. Self-regulation processes and health: the importance of optimism and goal adjustment. *J Pers* 2006;74:1721–47.
41. Harlow HF, Zimmermann RR. Affectional responses in the infant monkey. *Science* 1959;130:421–32.
42. Castenmiller JJ, West CE. Bioavailability and bioconversion of carotenoids. *Annu Rev Nutr* 1998;18:19–38.
43. Yeum KJ, Russell RM. Carotenoid bioavailability and bioconversion. *Annu Rev Nutr* 2002;22:483–504.
44. van Herpen-Broekmans WM, Klopping-Ketelaars IA, Bots ML, Kluft C, Princen H, Hendriks HF, Tijburg LB, van Poppel G, Kardinaal AF. Serum carotenoids and vitamins in relation to markers of endothelial function and inflammation. *Eur J Epidemiol* 2004;19:915–21.
45. van der Vaart H, Postma DS, Timens W, ten Hacken NH. Acute effects of cigarette smoke on inflammation and oxidative stress: a review. *Thorax* 2004;59:713–21.
46. Yanbaeva DG, Dentener MA, Creutzberg EC, Wesseling G, Wouters EF. Systemic effects of smoking. *Chest* 2007;131:1557–66.
47. Ikeda A, Schwartz J, Peters JL, Fang S, Spiro A, Sparrow D, Vokonas P, Kubzansky LD. Optimism in relation to inflammation and endothelial dysfunction in older men: the VA Normative Aging Study. *Psychosom Med* 2011;73:664–71.
48. Roy B, Diez-Roux AV, Seeman T, Ranjit N, Shea S, Cushman M. Association of optimism and pessimism with inflammation and hemostasis in the Multi-Ethnic Study of Atherosclerosis (MESA). *Psychosom Med* 2010;72:134–40.
49. Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C. Antioxidant supplements for prevention of mortality in healthy participants and patients with various diseases. *Cochrane Database Syst Rev* 2012;3: CD007176.
50. Ferrucci L, Perry JR, Matteini A, Perola M, Tanaka T, Silander K, Rice N, Melzer D, Murray A, Cluett C, Fried LP, Albanes D, Corsi AM, Cherubini A, Guralnik J, Bandinelli S, Singleton A, Virtamo J, Walston J, Semba RD, Frayling TM. Common variation in the beta-carotene 15,15'-monooxygenase 1 gene affects circulating levels of carotenoids: a genome-wide association study. *Am J Hum Genet* 2009;84:123–33.